

# Mapping Early Bone Marrow Lesion and Pain Development in Spontaneous and Load-induced mouse models of OA



Rebecca Norman¹, B. Javaheri¹, C. Chenu¹, N Sofat², AA. Pitsillides¹;
¹ Skeletal Biology Group, Comparative Biomedical Sciences, Royal Veterinary College, London, UK;
² Q• Œ Ở Á ÁQ △ &Œ ÞÁ ÁQ 〈 Šē ÁÐ ÁQ 〈 ` } Ē ÊÛdĚÕ^[ ¦\* ^q ÁŊ Ƹ^¦• æ Á Ƹ[ } å[ } ÊŠ[ } å[ } ÊVS

## What is osteoarthritis (OA)?

OA is a disease characterised by multiple soft and mineralised joint tissue changes, which have relationships that remain unresolved

Figure 1. Features of healthy (left) compared to OA joint (right)

## Aims and hypotheses

Characterise the temporal emergence of SCC and their spatial relationship to cartilage lesions in ageing STR/ort mice with increasing OA severity

Hypothesis: that SCC will increase in prevalence, number and volume with increasing age and OA grade, and will be closely aligned with the OA-prone sites of articular cartilage on the medial tibial plateau of STR/ort mouse joints

#### Methods and work flow

1. Left hindlimb selection (from archived samples):




Figure 3. Number of limbs per group

2. Knee joint imaged by µCT for SCC volume measurement

3

#### References

- [1] Yusuf, E. et al. (2011) Ann Rheum Dis, 70(1), 60-67.
- [2] Pouders, C. et al. (2008) Am J Roentgenol, 190(1), 17-21
- [3] Carrino, J. A. et al. (2006) Osteoarthr Cartil, 14(10), 1081-5. [4] Crema, M. D. et al. (2010) Radiol, 256(3), 855-62.